



E-ISSN: 2708-0064  
P-ISSN: 2708-0056  
JCRSI 2024; 6(1): 01-04  
[www.allcasereports.com](http://www.allcasereports.com)  
Received: 03-11-2023  
Accepted: 07-12-2023

**Dr. Shweta Garg**  
Consultant Pathologist,  
Consultant Histopathologist,  
Cytopathologist and  
Haematologist, Department of  
Pathology, Dist Hospital  
Chhatarpur & Saanvi  
Diagnostic, Madhya Pradesh,  
India

**Dr. Sanjana Robinson**  
Surgical Specialist, District  
Hospital Chhatarpur, Madhya  
Pradesh, India

## Axillary schwannoma: A rare occurrence

**Dr. Shweta Garg and Dr. Sanjana Robinson**

**DOI:** <https://doi.org/10.22271/27080056.2024.v6.i1a.68>

### Abstract

**Background:** Schwannomas are benign tumors arising from schwann cells. Most of the schwannomas occur in the head and neck region and in the flexor aspect of the limbs.

**Case presentation:** Herein we present a case of a 19 year old male who presented with a swelling in his left axilla for 3-4 months it was painless and not associated with any neurological deficit on the upper limbs.

The mass was excised and sent for histopathological examination. The diagnosis was confirmed by histopathology examination as schwannoma.

**Conclusion:** Schwannoma are benign nerve tumors of the nerve sheath. Schwannoma is rare in the axillary region. Surgical excision gives the good prognosis.

**Keywords:** Axillary swelling, schwannoma

### Introduction

Schwannomas are benign encapsulated perineural tumors arising from schwann cells of nerve sheath of peripheral or cranial nerves.

About 25% of all schwannomas occur in the head and neck region.

They rarely occur in the axillary region.

We report here one such case of axillary schwannoma which is rare and unusual in its site of occurrences.

**Case Report:** A 19 year old male patient presented to Government Hospital with swelling in the left axilla since 3-4 months, not associated with any pain or tingling sensation.



**Fig 1:** Site of axillary swelling

On examination solitary firm swelling of 4X3cm in its greatest dimension was situated in the left axilla.

Lymph nodes were not palpable.

There were no neurological deficits detected in the left upper limb.

All hematological investigations were within normal limits.

The patient underwent excision of the tumor that was found to be situated beneath the brachial plexus.

### Corresponding Author:

**Dr. Shweta Garg**  
Consultant Pathologist,  
Consultant Histopathologist,  
cytopathologist and  
Haematologist, Department of  
Pathology, Dist Hospital  
Chhatarpur & Saanvi  
Diagnostic, Madhya Pradesh,  
India

The patient did not have any neurological deficit after the procedure, except for the pain in the axilla. The patient was called for follow up after 1 month without any neurological deficit. Histopathological examination (HPE) showed encapsulated lesion.



**Fig 2:** Excised axillary swelling, nodular encapsulated mass measuring 4.0X3.0 cm, gross specimen and cut surface

Consisting of proliferating fibroblasts with scanty collagen and organoid arrangement of cells at places without cellular atypia, suggestive of schwannoma.

### Discussion

Schwannoma (Neurilemmoma) is a benign tumor arising from schwann cells.

Schwannomas most commonly affect people between the ages of 50 and 60 years, affecting both sexes equally, they rarely occur in children schwannomas are rare.

Schwannoma is a slow growing solitary firm, well circumscribed and encapsulated round or ovoid tumor.

The presence of large tumors as well as rare cases of multiple neoplasms has been reported.


An extracranial schwannoma may present as a solitary swelling anywhere in the body, the most common sites being the head and neck, flexor surfaces of the upper and lower limbs, posterior mediastinum in the thorax and on the trunk.

However a schwannoma in the axilla is of unusual occurrence.

Thus on account of their rare occurrence and involvement of unusual sites they pose a significant challenge to diagnosis and treatment.

Biopsy for HPE is an important tool for diagnosis.

On HPE schwannomas are found to contain a varying proportion of two different areas.



**Fig 3-11:** Histopathology, the H&E stained sections show Antoni A and Antoni B areas, with nuclear palisading, anuclear zone and Verocay bodies.

Antoni type A areas are highly cellular and contain closely packed spindle shaped cells forming palisades called verocay bodies.

Antoni type B areas are composed of loosely arranged

spindle shaped cells in a mucinous matrix the first choice of treatment for schwannoma is surgical excision of the tumor while sparing the nerve trunk to avoid neurological deficits.

## Conclusion

We have presented a rare case of axillary schwannoma. They pose a significant challenge to diagnosis on account of their rare occurrence in unusual sites. Because of the non-specific symptoms many cases can simply be missed and misdiagnosed. Hence a clinician has to bear in mind the possibility of a schwannoma in patients presenting with a swelling in the axilla. Surgical excision of these tumors is associated with good prognosis.

## Conflict of Interest

Not available

## Financial Support

Not available

## References

1. Zhou J, Man XY, Zheng M, Cai SQ. Multiple plexiform schwannoma of a finger. *Eur J Dermatol*. 2012;22:149-50. [PubMed] [Google Scholar]
2. Ackerman LV, Taylor FH. Neurogenous tumors within the thorax; a clinicopathological evaluation of forty-eight cases. *Cancer*. 1951;4:669-91. [PubMed] [Google Scholar]
3. Kehoe NJ, Reid RP, Semple JC. Solitary benign peripheral-nerve tumours. Review of 32 years' experience. *J Bone Joint Surg Br*. 1995;77:497-500. [PubMed] [Google Scholar]
4. Malizos K, Ioannou M, Kontogeorgakos V. Ancient schwannoma involving the median nerve: A case report and review of the literature. *Strategies Trauma Limb Reconstr*. 2013;8:63-6. [PMC free article] [PubMed] [Google Scholar]
5. Kang HJ, Shin SJ, Kang ES. Schwannomas of the upper extremity. *J Hand Surg Br*. 2000;25:604-7. [PubMed] [Google Scholar]
6. Ozdemir O, Ozsoy MH, Kurt C, Coskunol E, Calli I. Schwannomas of the hand and wrist: Long-term results and review of the literature. *J Orthop Surg (Hong Kong)* 2005;13:267-72. [PubMed] [Google Scholar]
7. Strickland JW, Steichen JB. Nerve tumors of the hand and forearm. *J Hand Surg Am*. 1977;2:285-91. [PubMed] [Google Scholar]
8. Ku HC, Yeh CW. Cervical schwannoma: A case report and eight years review. *J Laryngol Otol*. 2000;114:414-7. [PubMed] [Google Scholar]
9. Huang JH, Samadani U, Zager EL. Brachial plexus region tumors: A review of their history, classification, surgical management, and outcomes. *Neurosurg Q*. 2003;13:151-61. [Google Scholar]
10. Ergun T, Lakadamyali H, Derincek A, Tarhan NC, Ozturk A. Magnetic resonance imaging in the visualization of benign tumors and tumor-like lesions of hand and wrist. *Curr Probl Diagn Radiol*. 2010;39:1-16. [PubMed] [Google Scholar]
11. Traistaru R, Enachescu V, Manuc D, Gruia C, Ghilusi M. Multiple right schwannoma. *Rom J Morphol Embryol*. 2008;49:235-9. [PubMed] [Google Scholar]
12. Date R, Muramatsu K, Ihara K, Taguchi T. Advantages of intra-capsular micro-enucleation of schwannoma arising from extremities. *Acta Neurochir (Wien)* 2012;154:173-8. [PubMed] [Google Scholar]
13. Kwok K, Davis B, Kliot M. Resection of a benign brachial plexus nerve sheath tumor using intraoperative electrophysiological monitoring. *Neurosurgery*. 2007;60(42):316-20. [PubMed] [Google Scholar]
14. Shahid M, Ahmad SS, Vasenwala SM, *et al*. Schwannoma of the scrotum: Case report and review of the literature. *Korean J Urol* 2014; 55: 219–221.
15. An X, Zhu M, Zhang N, *et al*. Schwannoma of the vagina - a common tumor but a rare location: A case report. *Mol Clin Oncol* 2017; 7: 783–786.
16. Radojkovic M, Mihailovic D, Stojanovic M, *et al*. Large retroperitoneal schwannoma: A rare cause of chronic back pain. *J Int Med Res* 2018; 46: 3404–3410.
17. Zou F, Dai M, Zhang B, *et al*. Misdiagnosis of a giant intrapelvic schwannoma: A case report. *Oncol Lett* 2013; 6: 1646–1648.
18. Artico M, Cervoni L, Wierzbicki V, *et al*. Benign neural sheath tumors of major nerves: Characteristics in 119 surgical cases. *Acta Neurochir (Wien)*. 1997;139:1108–1116.
19. Yafit D, Horowitz G, Vital I, *et al*. An algorithm for treating extracranial head and neck schwannomas. *Eur Arch Otorhinolaryngol*. 2015;272:2035-2038.
20. Adani R, Baccarani A, Guidi E, *et al*. Schwannomas of the upper extremity: Diagnosis and treatment. *Chir Organi Mov*. 2008;92:85-88.
21. Châtillon CE, Guiot MC, Jacques L. Lipomatous, vascular and chondromatous benign tumors of the peripheral nerves. *Neurosurg Focus*. 2007;22:E18.
22. Panwar P, Kumar S, Singh S, *et al*. Giant abdominoperineal malignant schwannoma: An unusual presentation and surgical challenge. *Case Rep Urol*. 2015;2015:728062.
23. Kim SH, Kim NH, Kim KR, *et al*. Schwannoma in head and neck: Preoperative imaging study and intracapsular enucleation for functional nerve preservation. *Yonsei Med J*. 2010;51:938–942.
24. Hubert J, Landes G, Tardif M. Schwannoma of the median nerve. *J Plast Surg Hand Surg*. 2013;47:75-77.
25. Biswas D, Marnane CN, Mal R, *et al*. Extracranial head and neck schwannomas: A 10-year review. *Auris Nasus Larynx*. 2007;34:353-359.
26. Joshi R. Learning from eponyms: Jose Verocay and Verocay bodies, Antoni A and B areas, Nils Antoni and Schwannomas. *Indian Dermatol Online J*. 2012;3:215-219.
27. Nishio J, Mori S, Nabeshima K, *et al*. Successful enucleation of large multinodular/plexiform schwannoma of the foot and ankle. *Springerplus*. 2015;4:260.
28. Adani R, Tarallo L, Mugnai R, *et al*. Schwannomas of the upper extremity: Analysis of 34 cases. *Acta Neurochir (Wien)*. 2014;156:2325-2330.
29. Date R, Muramatsu K, Ihara K, *et al*. Advantages of intra-capsular micro-enucleation of schwannoma arising from extremities. *Acta Neurochir (Wien)*. 2012;154:173-178.
30. Uchida K, Nakajima H, Sato R, *et al*. Microsurgical intraneural extracapsular resection of neurinoma around the cervical neuroforamen: A technical note. *Minim Invasive Neurosurg*. 2009;52:271–274.
31. Ozdemir O, Ozsoy MH, Kurt C, *et al*. Schwannomas of the hand and wrist: long-term results and review of the literature. *J Orthop Surg (Hong Kong)*. 2005;13:267-



- 272.
32. Hirai T, Kobayashi H, Akiyama T, *et al.* Predictive factors for complications after surgical treatment for schwannomas of the extremities. *BMC Musculoskelet Disord.* 2019;20:166.
  33. Fujibuchi T, Miyawaki J, Kidani T, *et al.* Risk factors for neurological complications after operative treatment for schwannomas. *J Clin Neurosci.* 2017;46:136-140.
  34. Kim SM, Seo SW, Lee JY, *et al.* Surgical outcome of schwannomas arising from major peripheral nerves in the lower limb. *Int Orthop.* 2012;36:1721-1725.
  35. Oberle J, Kahamba J, Richter HP. Peripheral nerve Schwannomas-an analysis of 16 patients. *Acta Neurochir (Wien).* 1997;139:949-953.
  36. Takase K, Yamamoto K, Imakiire A. Clinical pathology and therapeutic results of neurilemmoma in the upper extremity. *J Orthop Surg (Hong Kong).* 2004;12:222-225.
  37. Sawada T, Sano M, Ogihara H, *et al.* The relationship between pre-operative symptoms, operative findings and postoperative complications in schwannomas. *J Hand Surg Br.* 2006;31:629-634.
  38. Kütahya H, Güleç A, Güzel Y, *et al.* Schwannoma of the median nerve at the wrist and palmar regions of the hand: A rare case report. *Case Rep Orthop.* 2013;2013:950106.

**How to Cite This Article**

Garg S, Robinson S. Axillary schwannoma: A rare occurrence. *Journal of Case Reports and Scientific Images.* 2024;6(1):01-04.

**Creative Commons (CC) License**

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.